

Resonating-valence-bond structure of Gutzwiller-projected superconducting wave functions

D. A. Ivanov

Institute of Theoretical Physics, École Polytechnique Fédérale de Lausanne (EPFL), 1015 Lausanne, Switzerland

Gutzwiller-projected (GP) wave functions have been widely used for describing spin-liquid physics in frustrated magnets and in high-temperature superconductors. Such wave functions are known to represent states of the resonating-valence-bond (RVB) type. In the present work I discuss the RVB structure of a GP singlet superconducting state with nodes in the spectrum. The resulting state for the undoped spin system may be described in terms of the “path integral” over loop coverings of the lattice, thus extending the known construction for RVB states. The problem of topological order in GP states may be reformulated in terms of the statistical behavior of loops. The simple example of the projected d -wave state on the square lattice demonstrates that the statistical behavior of loops is renormalized in a nontrivial manner by the projection.

Gutzwiller-projected (GP) wave functions have been often applied to describe the unconventional properties of high-temperature superconductors and the spin-liquid phase of frustrated spin-1/2 systems [1, 2, 3, 4]. Since the early days of their use for strongly correlated systems, GP wave functions have been known not only to provide a good variational ansatz, but also to represent the resonating-valence-bond (RVB) physics of the ground state. Because of their RVB structure, the GP wave functions merit deep investigation as independent objects, irrespective of the underlying physical Hamiltonian [5, 6].

It was shown in the original analysis of GP superconducting wave functions [1] that such wave functions correspond to the RVB states with singlet amplitudes a_{ij} given by the Fourier transform of

$$a(k) = \frac{u_k}{v_k}, \quad (1)$$

where u_k and v_k are the coherence factors of the BCS wave function before projection. For a fully gapped superconductor, this procedure produces amplitudes a_{ij} decaying exponentially with the site separation $|i - j|$, and the interpretation as a RVB state is straightforward. However, in the case of a BCS superconductor with nodes, including the commonly used $d_{x^2-y^2}$ state, this simple derivation produces values of $a(k)$ singular at the nodal points, and the resulting singlet amplitudes a_{ij} are long-ranged. Therefore, in the case of a superconductor with nodes, the conventional derivation cannot interpret the GP wave function as a RVB state.

In this note I resolve the problem of interpreting the physical content of the GP wave function by using an alternative representation in terms of the “loop-soup” path integral. In this construction, well-known for RVB states, the correlation functions are expressed as sums over all close-packed coverings of the lattice with closed loops [7, 8, 9]. In the conventional RVB loop construction, the statistical weights are determined by the products of singlet amplitudes along the loops. In my generalized derivation, the product of singlet amplitudes is replaced by the trace of the product of the BCS Green’s functions

along the loop. Thus the role of the RVB singlet amplitudes is played by the BCS Green’s functions (2×2 matrices). In the case of a superconductor with nodes, the BCS Green’s functions are only marginally local: they decay algebraically with distance. It is therefore clear that caution is required when applying the spin-liquid RVB scenario to high-temperature superconductors [10].

The paper is continued with a brief discussion of the relation of the loop-soup construction to the topological order predicted for RVB states [5, 6, 11, 12, 13]. The presence of topological order corresponds to disordered short-ranged loops.

I then employ the example of the undoped GP d -wave state on the square lattice to demonstrate the possibility of strong renormalization of the loop statistics by projection. The loop-length distribution may be characterized by spin-spin correlations which decay algebraically with distance. The power of the algebraic decay is non-universal and depends on the variational parameter of the wave function.

The derivation begins by defining the GP wave function. I consider a system consisting of spins 1/2 occupying a finite lattice with an even number of sites N . The GP wave function is constructed by using an auxiliary Hamiltonian of the BCS form,

$$H = \sum_{\{ij\}, \alpha} \left[t_{ij} \psi_{i\alpha}^\dagger \psi_{j\alpha} + \text{h.c.} \right] + \sum_{\{ij\}} \left[\Delta_{ij} (\psi_{i\uparrow}^\dagger \psi_{j\downarrow}^\dagger - \psi_{i\downarrow}^\dagger \psi_{j\uparrow}^\dagger) + \text{h.c.} \right] \quad (2)$$

(in this general form the hopping and pairing amplitudes between any pairs of sites are arbitrary). The Hamiltonian is spin-rotation invariant (it involves only singlet pairing), and its ground state Ψ_{BCS} is a spin singlet. The GP wave function Ψ_{GP} for the undoped system is constructed by projecting onto states with exactly one fermion per lattice site (Gutzwiller projection). The resulting wave function may be written as a wave function of spin variables $\Psi_{\text{GP}}(\{\sigma_i\})$, where the spins σ_i must contain $N/2$ up and $N/2$ down spins.

The Hamiltonian (2) contains more information than required for the construction of the wave function. First, the Gutzwiller projection selects states invariant with respect to SU(2) gauge rotations in the particle-hole space [14, 15]. Different Hamiltonians related by such SU(2) gauge symmetries therefore produce the same projected wave function Ψ_{GP} . Second, the Hamiltonian (2) contains information not only about its eigenfunctions, but also about the spectrum, which is not used in the construction of the wavefunction. To eliminate the latter redundancy, we shall use the equal-time Green's function (the “projector operator”) instead of the Hamiltonian.

To construct the equal-time Green's functions, we first define the $2N$ -dimensional vector space of fermionic operators

$$\gamma = \sum_i \left[u(i)\psi_{i\uparrow}^\dagger + v(i)\psi_{i\downarrow} \right] \quad (3)$$

(the sum is taken over the N lattice sites). This space has the Hermitian form defined by the anti-commutator $\{\gamma_1^\dagger, \gamma_2\}$, and the Hamiltonian (2) acts in this space as a Hermitian operator. The eigenvalue equations are the Bogoliubov–de Gennes equations

$$[H, \gamma] = E\gamma. \quad (4)$$

The spectrum of the eigenvalues consists of pairs of opposite energies $\pm E$: this is the consequence of the spin-rotational invariance of the Hamiltonian (2). For each solution (u, v) with energy E , the pair $(v^*, -u^*)$ gives a solution with energy $-E$. Thus there are N positive-energy solutions and N negative-energy solutions to Eq. (4).

Next we define the projector onto the negative-energy states,

$$G = \sum_{E_k < 0} |\gamma_k\rangle \langle \gamma_k|, \quad (5)$$

where $|\gamma_k\rangle$ are normalized eigenstates [solutions of (4)]. The operator G constructed in this way is a $2N \times 2N$ Hermitian matrix which may be considered as a single-particle projector onto the negative-energy states or, equivalently, as the matrix of equal-time Green's functions. It contains no information about the eigenvalues of (4), but only about the eigenfunctions. Thus the projector G contains less information than the original Hamiltonian, but remains sufficient for constructing the multi-particle ground-state wave function.

In real space, the operator G may be represented by the set of 2×2 matrices G_{ij} for each pair of lattice sites (i, j) . The hermiticity implies

$$G_{ij}^\dagger = G_{ji}. \quad (6)$$

The SU(2) gauge transformation acts on G by conjugation

$$G_{ij} \mapsto W_i^\dagger G_{ij} W_j, \quad (7)$$

where W_i are the SU(2) matrices of the gauge transformation.

From the completeness of the basis of all states γ_k and from the $E \rightarrow -E$ symmetry (i.e. from the spin-rotational invariance), we can derive the following identity for G_{ij} :

$$G_{ij} + J^\dagger G_{ij}^* J = \delta_{ij} \mathbf{1}, \quad J = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}. \quad (8)$$

Mathematically, this identity expresses the fact that $G - \frac{1}{2}$ belongs to the Lie algebra $\mathfrak{sp}(N)$ (see, for example, Ref. 16). One may verify that Eq. (8) is invariant with respect to SU(2) gauge rotations.

From the SU(2) invariance of Gutzwiller projection, it follows that the wave function depends only on the SU(2)-gauge-invariant properties of the Green's functions G_{ij} . For future application, we define the cyclic trace of the Green's functions

$$T_{i_1 \dots i_l} = \frac{1}{2} \text{Tr}(G_{i_1 i_2} G_{i_2 i_3} \dots G_{i_l i_1}), \quad (9)$$

which depends on the oriented closed loop $i_1 i_2 \dots i_l$. This type of cyclic trace appears below in the loop path integral.

To proceed with the loop construction, we first express the wave function in terms of the negative-energy states γ_k and further develop the loop path integral for ground-state expectation values.

The BCS ground state of the Hamiltonian (2) may be expressed as

$$\Psi_{\text{BCS}} = \prod_{E_k < 0} \gamma_k |\downarrow \dots \downarrow\rangle, \quad (10)$$

where $|\downarrow \dots \downarrow\rangle$ is the state with all lattice sites occupied by down spins, $|\downarrow \dots \downarrow\rangle = \prod_i \psi_{i\downarrow}^\dagger |0\rangle$, and the operators γ_k are the negative-energy eigenvectors of Eq. (4).

After projecting onto singly-occupied states, this wave function may be shown to yield for the GP wave function

$$\Psi_{\text{GP}}(\{\sigma_i\}) = \text{Det}_{k,i} [u_k(i)|v_k(i)], \quad (11)$$

where $[u_k(i)|v_k(i)]$ is the $N \times N$ matrix whose first $N/2$ columns are composed of the coefficients $u_k(i)$ while the last $N/2$ columns involve those of $v_k(i)$. The index k refers to all negative-energy states and the index i to all spin-up sites in the given spin configuration $\{\sigma_i\}$.

Alternatively, the same wave function may be written in the same form (11), but with the index i labeling spin-down sites. It follows from the rotation invariance that these two determinants must give the same wave function, to within an overall phase factor.

The expression for the “partition function” $\langle \Psi | \Psi \rangle$ is obtained by taking for the bra-vector the determinant (11) over the spin-up sites and for the ket-vector the same determinant over the spin-down sites. The product of the




FIG. 1: (a) An example of covering the lattice by loops in the “loop-soup” construction. (b) To each link of the loop, there corresponds a 2×2 matrix Green’s function G_{ij} . Each (oriented) loop is assigned the amplitude $-2T_{i_1 \dots i_l}$. The origin of the negative sign in the loop amplitude lies in the fermionic statistics in the GP construction. The vertices are labeled with up or down spins which alternate along the loop. The factor of two in the loop amplitude is due to the two possible spin labelings of the loop.

two determinants is then rewritten as the determinant of the product of the two matrices (taking the sum over the index k) to yield

$$|\Psi_{\text{GP}}(\{\sigma_i\})|^2 = \text{const Det}_{ij}[G_{ij}], \quad (12)$$

where i labels spin-up sites and j labels spin-down sites. The matrix $[G_{ij}]$ is the $N \times N$ matrix composed of 2×2 blocks G_{ij} . The constant prefactor does not depend on the spin configuration $\{\sigma_i\}$ and will be omitted in further calculations. The expression (12) is explicitly $\text{SU}(2)$ gauge-invariant.

The determinant (12) may be further expanded as a product of non-intersecting loops and summed over all possible spin configurations $\{\sigma_i\}$ to produce the partition function $\langle \Psi | \Psi \rangle$. Brief algebraic manipulations using the symmetry (8) give the loop path integral

$$\langle \Psi | \Psi \rangle = \sum_{\{C_n\}} \prod_n (-2T_{i_1 \dots i_{l_n}}), \quad (13)$$

where the sum is taken over all coverings by non-intersecting loops of even length, and $T_{i_1 \dots i_{l_n}}$ are the loop traces defined in Eq. (9), see Fig. 1. The sum (13) [and all loop sums below] is to be understood as that over *oriented* loops. From the symmetry (8) it follows that the traces $T_{i_1 \dots i_{l_n}}$ for even-length loops are purely real and do not depend on the orientation of the loop. However, it is important to preserve loop orientations in the definition of the sum (13), because they affect the multiplicities of length-two loops: every length-two loop has only one orientation and appears in the sum (13) once; all longer loops admit two orientations and appear twice.

This result has the form of a path integral in which different correlation functions may be computed. For example, the spin-spin correlation function $\langle S_z(i) S_z(j) \rangle$ may be written as

$$\langle S_z(i) S_z(j) \rangle = \frac{1}{\langle \Psi | \Psi \rangle} \sum_{\{C_n\}}^{i \leftrightarrow j} (-1)^{P_{ij}} \prod_n (-2T_{i_1 \dots i_{l_n}}), \quad (14)$$




FIG. 2: Defining topological sectors on a cylinder. The “reference line” (dashed) connects the two edges of the cylinder. (a) The “plus-minus” sectors in the loop path integral are generated by changing the sign of Green’s functions G_{ij} intersecting the reference line. (b) The “even-odd” sectors in the RVB states are selected as those with an even (odd, respectively) number of singlets intersecting the reference line.

where the sum is now taken only over loop coverings with the sites i and j belonging to the same loop, and $(-1)^{P_{ij}}$ takes values ± 1 depending on whether the number of loop links between i and j is even or odd.

The path integral (13) has the same form as the loop construction for RVB states. Most generally, the loop path integral may be written as

$$\langle \Psi | \Psi \rangle = \sum_{\{C_n\}} \prod_n A(C_n), \quad (15)$$

where $A(C_n)$ is an amplitude depending on the geometry of the loop C_n .

This general formulation includes many different RVB-type wave functions in a variety of systems. The GP wave functions in Anderson’s derivation [1] correspond to the loop amplitude

$$A(C_n) = -2a_{i_1 i_2} a_{i_2 i_3} \dots a_{i_l i_1}, \quad (16)$$

where a_{ij} are the singlet amplitudes defined from (1). Note the importance of the negative sign in the loop amplitudes (13) and (16). This sign arises from the fermionic statistics involved in the GP construction. Alternatively, for a spin system it is possible to construct a “bosonic” product of singlets which leads to a loop amplitude similar to Eq. (16), but *without* the negative sign [7, 8, 9].

Remarkably, the Rokhsar–Kivelson ground state of dimer models [11] may also be described with the same formalism (15): it is sufficient to set $A(C_n)$ equal to 1 for length-two loops on allowed dimer positions and to 0 otherwise.

The concept of the “loop soup” aids in visualizing the conditions required for the topological order proposed for RVB states [11, 12, 13]. As explained in Refs. 5, 6, the different topological sectors on a multiply connected domain (e.g. on a cylinder or a torus) may be accessed by imposing periodic or antiperiodic boundary conditions on the fermions in the Hamiltonian (2) along a topologically nontrivial contour. As a result, the periodic/antiperiodic boundary conditions produce two different GP wave functions Ψ_+ and Ψ_- . The topological order implies two dual




FIG. 3: **Main panel:** staggered spin correlations in the undoped projected d -wave state for different values of the variational parameter $\phi = 4 \arctan \tilde{\Delta}$. The Monte Carlo simulation was performed on the 24×24 lattice and involved averaging over 10^4 samples. The error bars are smaller than the symbol sizes. **Inset:** The exponent α as defined in Eq. (23) as a function of the variational parameter ϕ . Squares: values of α obtained from the spatial decay of spin correlations in the 24×24 system. Stars: values of α obtained from the size-dependence of the squared integrated staggered magnetization $M(L)$ defined in Eq. (24), see Fig. 4 below. The error bars are of the order of the symbol sizes. The arrow marks the position of the optimal value of the variational parameter ($\phi \approx 0.34\pi$) minimizing the variational energy of the Heisenberg Hamiltonian within the given class of wave functions [19, 20].

conditions

$$\langle \Psi_+ | X | \Psi_+ \rangle - \langle \Psi_- | X | \Psi_- \rangle \rightarrow 0, \quad (17)$$

$$\langle \Psi_+ | X | \Psi_- \rangle \rightarrow 0, \quad (18)$$

where the correlation functions must tend to zero sufficiently rapidly (e.g., exponentially) with increasing system size for any local operator X .

It will become clear below that the first of these conditions (17) may be viewed as an effective absence of infinite loops in the path integral (15), and may be related loosely to the short range of spin correlations. The second condition (18) corresponds to the absence of valence-bond crystallization in the RVB construction, and, in the general case (15), may in all probability be formulated as the absence of loop crystallization. Note, however, that the following discussion of those correspondences is not fully rigorous and some statements are indicated as conjectures.

I begin by considering the condition (17). Under the assumption that the Green's functions G_{ij} decay rapidly with distance, loops of large size may appear only




FIG. 4: The dependence of the fluctuating integrated staggered magnetization $M(L)$ defined in Eq. (24) on the linear system size L in the log-log scale. The values of α deduced from these data are plotted by star symbols in the inset of Fig. 3.

with many links of short length. In this case, the change of boundary conditions in Eq. (13) may be represented by changing the sign of the Green functions G_{ij} intersecting the reference line (Fig. 2). Consequently, the partition functions in the $\pm$ sectors may be written as

$$\langle \Psi_{\pm} | \Psi_{\pm} \rangle = \sum_{\{C_n\}} (\pm 1)^W \prod_n (-2T_{i_1 \dots i_n}), \quad (19)$$

where W is the total winding number of the loops. This definition may be extended directly to the general formalism (15) and to any expectation value of a local observable X ,

$$\langle \Psi_{\pm} | X | \Psi_{\pm} \rangle = \sum_{\{C_n\}} (\pm 1)^W X(\{C_n\}) \prod_n A(C_n). \quad (20)$$

Therefore the difference between the two topological sectors (17) contains only configurations with odd winding numbers. The absence of large loops then serves as a sufficient condition for the criterion (17) of topological order [I believe that in most situations this is also a necessary condition]. On the other hand, the absence of large loops also guarantees that spin correlations (14) are short-ranged, but is not a necessary condition. Therefore one may expect that in many cases the short range of spin correlations is related to the condition (17), but in this contribution I provide no rigorous derivation of such a relation.

We consider next the second condition (18). For the RVB wave functions represented as linear combinations of products of singlets, this condition corresponds to the absence of singlet ordering (e.g. the absence of a valence-bond crystal [17]). This can be shown by employing the

even-odd basis defined conveniently in the case of RVB or dimer states on a topologically non-trivial domain [13, 18]. Using the expression (20) for the definition of $\Psi_{\pm}$, the “plus-minus” basis is related to the “even-odd” basis by

$$\Psi_{\pm} = \Psi_e \pm \Psi_o, \quad (21)$$

where Ψ_o and Ψ_e are the states with even and odd numbers of intersections of the singlets (dimers) with the reference line (Fig. 2). The condition (18) may then be reformulated as

$$\langle \Psi_e | X | \Psi_e \rangle - \langle \Psi_o | X | \Psi_o \rangle \rightarrow 0 \quad (22)$$

which, in turn, can be interpreted as the absence of singlet (dimer) crystallization [18]. One may expect that for the more general loop construction (15) this condition also corresponds to the absence of loop crystallization. However I do not have a rigorous argument supporting this conjecture.

As discussed above, the first criterion of the topological order (17) involves the question of the effective loop size in the path-integral ensemble (15). This effective loop size cannot be inferred easily from the singlet amplitudes a_{ij} (in the case of the RVB construction) or from the Green’s functions G_{ij} . Numerical studies of spin correlations in “bosonic” RVB states [7] indicate that the effective loop size may be renormalized nontrivially by the constraint of fully-packed loops. Similar renormalizations of loop statistics may also be observed in the fermionic GP construction. I consider for illustration one of the simplest examples of a GP wave function, the undoped projected d -wave state on the square lattice. This state has been studied previously as an approximate ground state of the Heisenberg antiferromagnet [19, 20]. It does not possess long-ranged antiferromagnetic order, but only a power-law decay of antiferromagnetic correlations [6, 21]. The only dimensionless parameter in the wave function is the pairing strength $\tilde{\Delta} = \Delta/t$, and the wave function is invariant under $\tilde{\Delta} \rightarrow \tilde{\Delta}^{-1}$ (see, for example, Ref. 19). Equivalently, the same wave function may be defined as the projected normal staggered-flux state with a flux per plaquette of $\phi = 4 \arctan \tilde{\Delta}$ [14, 15, 22]. Numerical Monte Carlo simulations indicate that for all values of the parameter ϕ , the antiferromagnetic correlations obey the power law,

$$(-1)^{i-j} \langle S_z(i) S_z(j) \rangle \propto |i - j|^{-\alpha}, \quad (23)$$

see Fig. 3. One observes that the spin correlations decay very slowly (with $\alpha < 2$), which implies that the effective loop size in the loop path integral (15) is of the order of the system size. The condition (17) of the topological order is therefore expected to fail. Interestingly, in Ref. 5 this state was indeed classified as one without topological order, but on the basis of the failure of the *other*

condition (18) [the condition (17) was not considered for this state]. Note that in the current example (undoped projected d -wave state) the spin correlations (23) probe precisely the probability of the two sites i and j belonging to the same loop: the Green’s functions G_{ij} connect only sites of opposite antiferromagnetic sublattices, and therefore the term $(-1)^{P_{ij}}$ in (14) is equal to $(-1)^{i-j}$ independently of the loop configuration.

The actual value of the exponent α in the power law (23) remains a subject of controversy. One possible scenario is the universal value of α depending only on the symmetries of the wave function (for example, the authors of Ref. 23 propose the value $\alpha = 3/2$). However, my numerical results suggest a different possibility: a parameter-dependent non-universal exponent $\alpha(\phi)$. The values $\alpha(\phi)$ extracted from the spatial decay of spin correlations are shown in the inset of Fig. 3. To verify those results, I have also computed the square of the integrated staggered magnetization in the $L \times L$ system,

$$M(L) = \frac{1}{L^2} \left\langle \left[\sum_i (-1)^i S_z(i) \right]^2 \right\rangle. \quad (24)$$

The dependence of $M(L)$ on the linear system size L is plotted in Fig. 4 for three values of ϕ . A simple scaling argument predicts $M(L) \propto L^{2-\alpha}$. The values of $\alpha(\phi)$ deduced from $M(L)$ are plotted by star symbols in the inset of Fig. 3. While those results are consistent with a ϕ -dependent value of α , they do not constitute a definite proof: a more detailed analysis of finite-size effects or an analytic argument is needed to settle the issue.

To summarize, the GP superconducting wave functions admit a loop description generalizing the conventional loop construction for RVB states. The role of the singlet amplitudes is played by the equal-time BCS Green’s functions. The properties of topological order and fractionalization are related to the correlations of the loops, but these cannot be inferred simply from the unprojected Green’s functions involved. Nevertheless, the loop formulation may be helpful in visualizing the RVB properties of the GP wave functions, and possibly for developing appropriate analytic approximations.

The author thanks G. Jackeli and B. Normand for helpful comments on the manuscript.

-
- [1] P. W. Anderson, Science **235**, 1196 (1987).
 - [2] C. Gros, Phys. Rev. B, **38**, 931 (1988); Ann. Phys. **189**, 53 (1989).
 - [3] L. Capriotti, F. Becca, A. Parola, and S. Sorella, Phys. Rev. Lett. **87**, 097201 (2001).
 - [4] P. W. Anderson et al., J. Phys. Cond. Matt. **16**, R755 (2004) [cond-mat/0311467].
 - [5] D. A. Ivanov and T. Senthil, Phys. Rev. B **66**, 115111 (2002).

- [6] A. Paramekanti, M. Randeria, N. Trivedi, Phys. Rev. B **71**, 094421 (2005).
- [7] S. Liang, B. Douçot, and P. W. Anderson, Phys. Rev. Lett. **61**, 365 (1988)
- [8] B. Sutherland, Phys. Rev. B **37**, 3786 (1988).
- [9] M. Kohmoto and Y. Shapir, Phys. Rev. B **37**, 9439 (1988).
- [10] T. Senthil and M. P. A. Fisher, Phys. Rev. B **63**, 134521 (2001); Phys. Rev. Lett. **86**, 292 (2001).
- [11] D. S. Rokhsar and S. A. Kivelson, Phys. Rev. Lett. **61**, 2376 (1988).
- [12] N. Read and B. Chakraborty, Phys. Rev. B **40**, 7133 (1989).
- [13] N. Bonesteel, Phys. Rev. B **40**, 8954 (1989).
- [14] I. Affleck, Z. Zou, T. Hsu, and P. W. Anderson, Phys. Rev. B **38**, 745 (1988).
- [15] F. C. Zhang, C. Gros, T. M. Rice, and H. Shiba, Supercond. Sci. Technol. **1**, 36 (1988).
- [16] A. Altland and M. R. Zirnbauer Phys. Rev. B **55**, 1142 (1997).
- [17] C. Lhuillier, cond-mat/0502464 and references therein.
- [18] A. Ioselevich, D. A. Ivanov, and M. V. Feigelman, Phys. Rev. B **66**, 174405 (2002).
- [19] H. Yokoyama and H. Shiba, J. Phys. Soc. Jpn. **57**, 2482 (1988); H. Yokoyama and M. Ogata, J. Phys. Soc. Jpn. **65**, 3615 (1996).
- [20] D. V. Dmitriev, V. Ya. Krivnov, V. N. Likhachev, and A. A. Ovchinnikov, Fiz. Tv. Tela. **38**, 397 (1996) [Phys. Solid State **38**, 219 (1996)].
- [21] D. A. Ivanov, Ph.D. thesis, M.I.T. 1999.
- [22] D. A. Ivanov and P. A. Lee, Phys. Rev. B **68**, 132501 (2003).
- [23] A. Paramekanti, M. Randeria, and N. Trivedi, Phys. Rev. B **70**, 054504 (2004).